

IPEF, n.23, p.29-36, abr.1983

## **Biomass and Nutrient Estimates in Short Rotation Intensively Cultured Plantation of *Eucalyptus grandis*<sup>1</sup>**

POGGIANI, F. and COUTO, H. T. Z.  
ESALQ -USP, Depto. de Silvicultura - 13.400 - Piracicaba -SP

SUITER FILHO, W.  
Cia. Agro-Florestal Santa Bárbara - Av. Brasil, 709 - 30.000 - Belo Horizonte - MG

**ABSTRACT** - Twenty four trees, including all diameters of an *Eucalyptus grandis* stand (2.5 years old - 5333 trees/ha) were weighted by component parts (leaves, limbs and stems). Samples were oven-dried to determine moisture content and to establish: leaves, limbs, crown, and stem weight-relationships. Also macro and microelements of each tree component were analyzed to estimate the nutrient contents removed by bole wood, harvest or complete trees utilization. Biomass distribution among the components of the stand is about 9% leaves, 7% limbs and 83% stems. However nutrients content in the stand biomass are about 37% in the leaves, 10% in the limbs and 53% in the stems. Because "cerrado" soil (Savanna) is very low in available form of nutrients, mainly P and K, it is suggested to avoid the removal of leaves, limbs and bark from the site.

**RESUMO** - Vinte e quatro árvores, incluindo todas as classes de diâmetro de um talhão de *Eucalyptus grandis* (com 2,5 anos - 5333 árvores/ha) foram selecionadas, cortadas e seus componentes (folhas, galhos e tronco) pesados. Amostras dos componentes foram secos em estufa para determinação da umidade e para estabelecer as relações entre as folhas, os galhos e o tronco. Foram, também estimados os nutrientes removidos pelos componentes de estande. Observou-se que 9% da biomassa é contida nas folhas, 7% nos galhos e 83% nos troncos. Todavia, 37% dos nutrientes são contidos nas folhas, 10% nos galhos e 53% nos troncos. Face à baixa fertilidade dos solos de cerrado, principalmente com relação ao P e K, seria oportuno evitar a remoção das folhas, galhos e cascas do "site".

### **Introduction**

Escalating cost of oil and energy shortage has forced wood users to consider maximizing total tree use. In fact, crown biomass and tree bark may become sources of energy. Therefore, intensive management of forest plantations will increase significantly biomass production.

In this area several papers have been published in Europe and North America (HANSEN & BAKER, 1979), however only a few studies were conducted in Brazil. Preliminary data have been published by POGGIANI, COUTO & SIMÕES (1979) regarding biomass production and nutrient accumulation in forest stand of *Eucalyptus grandis* planted in the State of São Paulo.

---

<sup>1</sup> Presented at the IUFRO Fast-Growing Trees Symposium, Águas de São Pedro - S. Paulo, August 1980.

According to HANSEN & BAKER (1979), intensive management of plantations may significantly increase biomass production (3 to 5 times), however it increases also nutrient removal from the site and a strong fertilization must become an integral part of such management.

POGGIANI (1980) reports some data regarding (*Eucalyptus* and *Pinus* plantations in the states of São Paulo and Minas Gerais and believes to be primordial to evaluate silvicultural and ecological consequences of intensively cultured plantations and short rotations in tropical areas. Cerrado soils (Brazilian Savannas) have in general low fertility, mainly in P and K, and high concentrations of iron, manganese and aluminium.

This paper discusses the distribution of biomass and nutrients in a stand of *Eucalyptus grandis* (2.5 years old). Equations predicting the dry biomass of the total tree and its components are presented. Nutrient removal by conventional bole-wood harvest and complete tree utilization are also discussed.

### **Site and Stand Description**

Seedlings of *Eucalyptus grandis* (Hill ex- maiden) were produced from seeds provenient from Coff's Harbour (Australia) and planted in November 1976. The area of this experiment, located in Bom Despacho (Minas Gerais) presents the typical climate of Brazilian savannas: annual mean temperature around 19.5° C with occasional frost in the winter; annual rainfall about 1400 mm, with 80% of the total rain-fall concentrated on the hot-wet season (October-March).

Each seedling planted in the initial spacing of 1.0 by 1.5 m was fertilized with 150 g of N-P-K (10:28:6) and B and Zn.

The planting area was previously occupied by a kind of savanna vegetation (Cerrado) and the soil is a clay latosol very deep and with a low content of available nutrients such as phosphorus 33 kg/ha, potassium 99 kg/ha, calcium 312 kg/ha, and magnesium 220 kg/ha (0-120 cm depth). The survival rate was around 80% at the end of the experiment.

### **Field and Laboratory Methods**

A random sample of 150 trees was selected from the 2.5 years old plantation of *Eucalyptus grandis* D.B.H. and the total height data of each tree were determined and the sample was divided in nine diameter classes. Two to four trees from each D.B.H. class from 2.0 to 11.0 cm (total 24 trees) were selected, felled and limbed. Each component of the tree (leaves, limbs and stem) was separated and weighted in the field.

Three samples (about 150 g) of leaves and small sections of the limbs collected from the median part of the crown and a disk collected from the median part of the stem were weighted and sealed in polyethylene bags for laboratory analysis. All the samples were oven-dried at 80°C until constant weight. After this moisture content of the different components were determined and the dry weight of each tree calculated. Weighted values for moisture content in the samples were used to convert component green weight to oven-dry weight.

After oven-drying leaves, limbs and stem samples (sections of the disks including wood and bark taken at half of the tree total height) were ground in a Wiley mill and passed through a 20 mesh screen. The chemical analyses were performed according to SARRUGE


& HAAG (1974). Nitrogen by the micro-Kjeldahl method, phosphorus by the vanadomolybdate method and all other nutrients were determined from the acid solution by atomic absorption spectrophotometer.

In order to estimate biomass content of the *E. grandis* stand regression models were determined and the parameters were estimated using the S.A.S. (Statistical Analysis System) computer package.

For choosing the best model to estimate dry biomass components of the tree a step-wise procedure was performed using as independent variables D.B.H. and total height, and their transformed values.

**TABLE 1: Biomass distribution (Kg/ha) of a *Eucalyptus grandis* stand for different diameter classes.**

CLASSES D.B.H.	TREES /ha	LEAVES kg/ha	LIMBS kg/ha	CROWS kg/ha	STEMS kg/ha	TOTAL TREES kg/ha
1) 2.1-3.0	261	20.3	106.8	121.8	180.2	307.3
2) 3.1-4.0	213	36.4	76.4	112.8	339.0	451.8
3) 4.1-5.0	480	209.3	208.3	417.6	1,722.7	2,140.3
4) 5.1-6.0	747	564.0	404.1	968.1	5,354.5	6,322.6
5) 6.1-7.0	1,013	1,161.0	772.9	1,939.9	11,108.5	13,048.4
6) 7.1-8.0	1,013	1,661.3	1,170.0	2,831.3	15,833.2	18,664.5
7) 8.1-9.0	907	1,892.0	1,454.0	3,346.0	18,038.4	21,384.4
8) 9.1-10.0	480	1,350.2	1,039.2	2,389.4	12,885.6	15,275.0
9) 10.1-11.0	213	117.4	553.4	1,270.8	6,849.2	8,120.0
TOTAL	5,333	7,617.9	5,785.1	13,403.0	72,311.3	85,714.3
%	(100%)	(8.9%)	(6.7%)	(15.6%)	(84.4%)	(100.0%)





$$Y = - 0.3145 + 0.0197 D^2 .H \quad R^2 = 0.99^{**}$$

Table 1 shows the dry biomass data for all classes of diameter and the total biomass produced by the stand of *E. grandis*.

If the complete tree utilization is considered, the total biomass amounts 85,7 tons per hectare. Conventional bole wood harvest of the stand would provide 72.3 tons of wood biomass (About 29 tons/ha per year).

These data compared with several data presented by HANSEN & BAKER (1979) including biomass production per year of some fast growing trees species in temperate zones shows clearly the strong potential of Eucalyptus for wood production in the tropics.

Spacing has also a strong influence in biomass produced by short rotations. Very closed spacing, according to BELANGER & PEPPER (1978) increased biomass production in sycamore plantations, however the rate of survival became lower in very closed spacing (0.4 m<sup>2</sup>).

Presently several experiments with closed spacing and short rotations are conducted by the Department of Silviculture of São Paulo University (Piracicaba). Preliminary data show the spacing 1.0 by 1.0 as the best for *Eucalyptus saligna* and *Eucalyptus alba* plantations, one year old, with a survival rate of 94%.

Anyway, short rotation increases strongly the biomass production. Comparing the biomass produced in this experiment with an 7 years old stand of *E. grandis* planted in the conventional spacing-3.0 by 2.0 m -(BELLOTE; 1979 and POGGIANI, 1980) it is possible to conclude that short rotations have twice the potential for wood production than usual rotations for *Eucalyptus* trees in Brazil.

In short rotations biomass distribution is also different comparing with conventional plantations. According to BELLOTE (1979) an *Eucalyptus grandis* stand - 2 years old and 3.0 by 2.0 m spacing - presented the following proportion for biomass distribution in the different components of the trees: 68.3% in the stem, 12.2% in the leaves and 19.5% in the limbs. The present experiment found for *E. grandis* (2.5 years old plantation) 83.2% of the biomass in the stem, 8.9% in the leaves and 7.9% in the limbs. It indicates more closed spacing in short rotations increase stem biomass. However more experiment must be carry out in this sense.

**TABLE 2: Biomass distribution on the average tree for different diameter classes**

CLASSES	D.B.H. cm	LEAVES Kg	LINES Kg	CROWN Kg	STEM Kg	TOTAL TREE Kg
1	2.1-3.0	0.016 (6.6%)	0.400 (34.6%)	0.476 (41.3%)	0.675 (58.7%)	1.151 (100.0%)
2	3.1-4.0	0.171 (8.0%)	0.357 (16.8%)	0.528 (24.8%)	1.594 (15.2%)	2.122 (100.0%)
3	4.1-5.0	0.436 (9.8%)	0.434 (9.7%)	0.870 (19.5%)	3.589 (80.5%)	4.459 (100.0%)
4	5.1-6.0	0.755 (8.9%)	0.541 (6.4%)	1.296 (15.3%)	7.168 (84.7%)	6.464 (100.0%)
5	6.1-7.0	1.152 (8.9%)	0.763 (5.9%)	1.915 (14.8%)	10.966 (85.2%)	12.881 (100.0%)
6	7.1-8.0	1.640 (8.9%)	1.155 (6.3%)	2.795 (15.2%)	15.630 (84.8%)	18.425 (100.0%)
7	8.1-9.0	2.085 (8.8%)	1.604 (6.8%)	3.689 (15.6%)	19.888 (84.4%)	23.577 (100.0%)
8	9.1-10.0	2.813 (8.8%)	2.165 (6.8%)	4.978 (15.6%)	26.845 (84.4%)	31.823 (100.0%)
9	10.1-11.0	3.368 (8.8%)	2.598 (6.8%)	5.966 (15.6%)	32.156 (84.4%)	38.122 (100.0%)

In the Table 2 are shown the dry biomass data per tree, for each class of diameter. Crown weight of the lower class is 41.3% of the total tree weight, all although in the higher class crown weight is only 15.6% of the tree.

This remark will be discussed in the item 3.4

#### Nutrient concentration in the tree components

Means and standard errors of nutrient concentration are included in table-3. For all the components nutrient concentrations in the biomass are as follows:

$N > K > Ca > Mg > p > Mn > Fe > Zn > Cu$

Nutrients concentration is for all the elements higher in the leaves than in the stem: 12 times for nitrogen, 4:5 times for phosphorus, 2.8 times for potassium, 6.5 times for calcium and 9.5 times for magnesium.

In general, nutrient concentrations found in this experiment are similar to the concentrations presented by HAAG et alii (1976) and BELLOTE (1979) for *E.grandis* plantations in "Cerrado" soil however they are lower than concentrations presented by LUBRANO (1967) for *Eucalyptus viminalis* planted in Italy.

Certainly N.P.K fertilization in the planting (150 g -10:28:6) makes available a regular supply of N and K, but P is strongly fixed by aluminium in the soil (pH 4.6), that is also very poor in calcium. Also high levels of Mn and Fe were observed in the leaves due to soil acidity, however no symptoms of toxicity were evident in the trees.

#### Nutrient content in the biomass of the stand

Recently in Brazil some foresters believe to be suitable whole tree utilization, mainly for energy supplies. In this sense, several researches are conducted in Brazil and in other countries. A critical evaluation about this problem was published by SWITZER, NELSON & HINESLEY (1978) but only for temperate zone. It is necessary to point out the problem is more complicated in the fragile ecosystems of the tropics, because the ecological and economical implications.

Table 4 presents the data on biomass and nutrients distribution in the stand.

**TABLE 3: Mean and standard error of nutrient concentrations in the trees components of *Eucalyptus grandis* (2.5 years old)**

COMPONENTES	ELEMENTS								
	N	P	K %	Ca	Mg	Fe	Cu ppm	Mn	Zn
LEAVES	1.850	0.110	0.650	0.590	0.200	296.0	4.0	953.0	12.5
S. Er.	0.020	0.012	0.050	0.060	0.014	33.0	0.2	102.0	0.6
LIMBS	0.460	0.040	0.370	0.260	0.040	93.0	3.6	396.0	5.7
S. Er.	0.017	0.003	0.030	0.023	0.007	10.1	0.1	26.3	0.4
STENS	0.150	0.024	0.230	0.090	0.021	38.6	1.5	185.0	5.3
S. Er	0.009	0.003	0.007	0.006	0.002	2.4	0.1	13.8	0.3

**TABLE 4: Biomass and nutrients content (Kg/ha) in an *Eucalyptus grandis* stand, 2.5 years old (5,333 trees/ha)**

COMPONENTS	BIOMASS	N	P	K	Ca	Mg	Fe	Cu	Mn	Zn	TOTAL NUTR.
TOTAL TREE	85,714.3	275.9	28.0	237.2	125.0	32.7	5.5	0.16	22.8	0.52	727.7
LEAVES	7,617.9	140.9	8.4	49.5	44.9	15.2	2.2	0.03	7.2	0.09	268.4
LIMBS	5,785.1	26.6	2.3	21.4	15.0	2.3	0.5	0.02	2.3	0.03	70.4
CROWN	13,403.0	167.5	10.7	70.9	59.9	17.5	2.7	0.05	9.5	0.12	338.8
STEM	72,311.3	108.4	17.3	166.3	65.1	15.2	2.8	0.11	13.3	0.40	388.9

Leaves and limbs constitute 15.6% of the total biomass, but contain 46,6% of the nutrients (338.8 kg/ha). Stems content is 388.9 Kg/ha. Nitrogen is the nutrient highly accumulated in the crown and potassium in the stem.

According to POGGIANI (1980) total biomass production of a stand of *E. grandis* in short rotation intensively cultured plantation might reach, in an eight years period, twice the production of a conventional rotation stand (3.0 by 2.0 m spacing). However the short rotation removes three times more nutrients than conventional plantations.

With the purpose to reduce the strong nutrient remotion by short rotations other alternatives of management and exploitation must be developed. Moreover new species provenances and clones with a higher "utilization efficiency" for closed spacing and short rotations must be introduced in silviculture practices.

According to HANSEN & BAKER(1979) we use ""utilization efficiency" to express the weight of biomass produced per unit weight of nutrients.

Fertilization versus nutrients removed by trees harvest

As was shown before, the "Cerrado" soil where this experience was laid down is poor in available nutrients (P-33 Kg/ha, K-99Kg/ha, Ca312 Kg/ha and Mg -220Kg/ha.). Comparing nutrient content of above ground biomass of Eucalyptus stand, as show in Table 5, with soil nutrients (0-120 cm depth), it can be seen that this soil does not support for a long time a forest plantation intensively exploited.

In fact, nutrient, content in the soil is almost equivalent to nutrient content in the biomass of the Eucalyptus stand. Mainly for potassium, phosphorus and calcium will be necessary to pay a supplementary attention. In such case is necessary to supply the nutrients removed by harvesting with a suitable fertilization to maintain the stand productivity.

In this experiment each tree received as fertilizer 150g of N-P-K (10:28:6) corresponding to 15g of nitrogen, 18.3g of phosphorus and 7.5g of potassium.

Probably, nitrogen is also fixed from the air by symbiotic and asymbiotic process, however a large rate of P is fixed by aluminum in the soil and potassium is accumulated in the plant tissue. Low content of potassium in the soil can become very critical for the next rotations. Calcium is also a scarce element in the soil and must be supplied by fertilization. This calcium scarcity is evident if we compare its concentration in the trees tissues with some data reported by LUBRANO (1967) for *Eucalyptus viminalis* in Italy. Probably soil liming before planting may result in more available nutrients for root systems and better economy in fertilizers.

### **Conclusion**

Some considerations are also essential to preserve nutrients in the site. Leaves and crown exploitation seem to be not convenient under an ecological and silvicultural point of view, mainly for dominated trees that present a large proportion of crown biomass with high rate of nutrients.

Therefore, to mitigate the loss of soil fertility in tropical zones would be suggested to avoid crown exploitation mainly in short rotation regime. Another suggestion would be to encourage research for species and clones selection with high "utilization efficiency" in order to improve biomass production with a lower nutrient remotion from the site.

According to BAKER & COOPER (1978) and BOWERSOX & WARD (1976) considerable variation in nutrient concentration has been noted in eastern cottonwood and in young hybrid poplars growing in closed spacing.

Fertilization studies are also primordial to develop more appropriated know-how for tropical soils of the Brazilian savannas.

**TABLE 5: Average of biomass (Kg) and nutrients content (g) per tree of *Eucalyptus grandis*, 2.5 years old**

COMPONENTS	BIOMASS	N	P	K	Ca	Mg	Fe	Cu	Mn	Zn	TOTAL NUTR.
g/trees											
TOTAL TREE	16.071 (100%)	51.7	5.2	44.5	23.4	6.0	1.0	0.038	4.3	0.054	136.2 (100%)
LEAVES	1.428 (8.9%)	26.4	1.6	9.3	8.4	2.8	0.4	0.005	1.4	0.018	50.3 (36.9%)
LIMBS	1.084 (6.7%)	5.0	0.4	4.0	2.8	0.4	0.1	0.004	0.4	0.006	13.1 (9.6%)
CROWN	2.512 (15.6%)	31.4	2.0	13.3	11.2	3.2	0.5	0.009	1.8	0.024	63.4 (46.5%)
STEM	13.559 (84.4%)	20.3	3.2	31.2	12.2	2.8	0.5	0.020	2.5	0.070	72.8 (53.5%)

### References

- BELANGER. R.P. & PEPPER, W.D. - Seedling density influences the early growth of planted sycamore. *Forest Science*, Madison. 24 (4):193-6. 1978.
- BELLOTE, A.F.J. Concentração, acumulação e exportação de nutrientes pelo *Eucalyptus grandis* (Hill, ex-Maiden) em função da idade Piracicaba. 1979. (Tese- Mestrado-ESALQ/USP).
- BLACKMON. B.G.; BAKER. J.B. & COOPER. R.T. - Biomass and nutrient accumulation in the geographic sources of eastern cottonwood. *Agronomy Abstracts*. 1978 Annual Meeting p.186.
- BOWERSOX. J.W. & WARD. W.W. - Growth and yield of close-spaced young hybrid poplars. *Forest Science*. Madison. 22(4): 449-59,1976.
- HAAG. H.P. et alii. - Análise foliar em cinco espécies de eucaliptos. *IPEF*, Piracicaba. (13):99-116.1976.
- HANSEN, E.A. & BAKER, J.B. - Biomass and nutrient removal in short rotation intensively cultured plantations. *Annual meeting North American Poplars Council*. Thompsonville, 1979 p.130-51.
- LUBANO, L. - Ricerche sulle esigenze nutritive di alcune speci di eucalitti. *Pubblicazioni Centro di Sperimentazione Agricola e Forestal*, Roma, 11:1-15 1972.
- POGGIANI, F. - Florestas para fins energéticos e ciclagem de nutrientes. *Série técnica, IPEF*, Piracicaba, 1(2): D1-D11, 1980.

POGGIANI, F.; COUTO, H.T.Z. do & Simões, J.W. - Aspectos ecológicos das minirotações e do aproveitamento dos resíduos florestais. *Circular Técnica IPEF*, Piracicaba, (74): 1-7, 1979.

SARRUGE, J.R. & HAAG, H.P. - *Análise química das plantas*. Piracicaba, ESALQ/USP 1974. 56p.

SWITZER, G.L.; NELSON, L.E. & HINESLEY, L.E. - Effects of utilization on nutrient regimes and site productivity. In: McMILLIN, C.W. - *Complete tree utilization of southern pine*. Madison, Forest Products Research Society, 1978 p.91-102 1978.